

sine.¹⁰ Cyclization of **4** to a tetrahydrobenzazepine **5** could be followed by oxidative fission of one aromatic ring in a manner analogous to the degradation of aromatic rings observed in a variety of living systems.¹¹ Two successive cyclizations transform **5** into a tetracyclic compound, **6**. Rearrangement of **6** could then give desmethylcephalotaxinone (**7**), which has been isolated from *Cephalotaxus* plants.¹² This hypothesis predicts that cephalotaxine should be labeled as shown in **7**. Experiments are in progress to determine if this prediction is correct.

Acknowledgment. We wish to thank Dr. R. G. Powell for gifts of alkaloids and Dr. Robert F. Perdue for dried plant material. Financial support from the National Institutes of Health (GM-19220) is gratefully acknowledged as is support from the National Science Foundation (GU 3852, GP 37156) which allowed purchase of a Bruker WH-90 NMR spectrometer used in this work.

References and Notes

- (1) W. W. Paudler, G. I. Kerley, and J. B. McKay, *J. Org. Chem.*, **28**, 2194 (1963); R. G. Powell, D. Weisleder, C. R. Smith, Jr., and I. A. Wolff, *Tetrahedron Lett.*, 4081 (1969); D. J. Abraham, R. D. Rosenstein, and E. L. McGandy, *ibid.*, 4085 (1969); S. K. Arora, R. B. Bates, R. A. Grady, and R. G. Powell, *J. Org. Chem.*, **39**, 1269 (1974).
- (2) R. G. Powell, D. Weisleder, C. R. Smith, Jr., and W. K. Rohwedder, *Tetrahedron Lett.*, 815 (1970); K. L. Mikolajczak, R. G. Powell, and C. R. Smith, Jr., *Tetrahedron*, **28**, 1995 (1972).
- (3) R. G. Powell, *Phytochemistry*, **11**, 1467 (1972).
- (4) I. D. Spenser, *Compr. Biochem.*, **20**, 231 (1968).
- (5) Feeding experiments were carried out in a Lab-line Biotronette Mark III environmental chamber.
- (6) Incorporation figures are based on the quantity of L-tyrosine fed.
- (7) E. Leete, *J. Am. Chem. Soc.*, **85**, 473 (1963).
- (8) All new compounds gave spectral data consistent with the assigned structures as well as satisfactory analytical figures via combustion analysis or high-resolution mass spectrometry.
- (9) J. M. Schwab and R. J. Parry, unpublished work.
- (10) E. Leete and A. Ahmad, *J. Am. Chem. Soc.*, **88**, 4722 (1966).
- (11) G. H. N. Towers and P. V. Subbo Rao, *Recent Adv. Phytochem.*, **4**, 1 (1972).
- (12) R. G. Powell and K. L. Mikolajczak, *Phytochemistry*, **12**, 2987 (1973).

Ronald J. Parry,* John M. Schwab

Department of Chemistry, Brandeis University
Waltham, Massachusetts 02154

Received December 14, 1974

Laser Separation of Chlorine Isotopes. The Photochemical Reaction of Electronically Excited Iodine Monochloride with Halogenated Olefins

Sir:

We report here the photochemical separation of ³⁵Cl and ³⁷Cl when a mixture of ICl and the scavengers *trans*-ClHC=CHCl and 1,2-dibromoethylene (*cis*, *trans* mixture) is irradiated by a CW tunable dye laser that selectively excites I³⁷Cl. The scavenger has the property that it does not react with ground state ICl but does react with ICl in the excited A ³Π₁ state. Thus, from *trans*-ClHC=CHCl, both the photoproduct *cis*-ClHC=CHCl and the starting material *trans*-ClHC=CHCl show ³⁷Cl-³⁵Cl exchange. The latter demonstrates for the first time laser-controlled isotope interchange. In the case of BrHC=CHBr, all photoproducts are ³⁷Cl enriched; in particular, the *trans*-ClHC=CHCl photoproduct has a ³⁵Cl:³⁷Cl ratio of 2:1 compared to 3:1 for naturally occurring *trans*-ClHC=CHCl. This technique of selective laser excitation and subsequent isotope labeling provides us with a new means of following organic gas-phase photochemical reactions with state selection of the reactants.




Figure 1. Potential energy curves for the X, A, and B states of ICl, taken from ref 1b with minor modification.




Figure 2. Low-resolution visible absorption spectrum of ICl taken from 6000 to 6100 Å. Note that the separation between the I³⁵Cl and I³⁷Cl bandheads are about 15 Å in this spectral region.

The visible absorption spectrum of I³⁵Cl and I³⁷Cl has been extensively studied,¹ and the relevant potential energy curves are shown in Figure 1. By tuning the CW dye laser to the (18, 0) bandhead of the I³⁷Cl A-X system at 6053 Å (see Figure 2), we can be assured that no photodissociation or photopredissociation occurs because the *v'* = 18 level of the A state lies ~784 cm⁻¹ below the X state dissociation limit. Our laser has an average power of 10 mW and a relatively wide band width (3 Å); this permits preferential excitation of the I³⁷Cl molecule, although some I³⁵Cl is also excited.

A 1:1 mixture of ICl and *trans*-ClHC=CHCl (10 Torr each) is placed inside a Pyrex tube 1 m long and 5 cm diameter, and exposed for 3 hr. In the single-pass reaction cell about one-third of the laser beam (5-mm spot size) is absorbed. Then excess C₂H₄ is added to trap unreacted ICl. The products are separated by gas-liquid phase chromatography (GC). The presence of an extra peak in the GC spectrum is analyzed to be the photoproduct *cis*-ClHC=CHCl.

Isotope analysis of the products is accomplished using a Nier-type mass spectrometer having an accuracy in the ratio of the mass peaks, *P*, of four parts per thousand. For starting material, *trans*-ClHC=CHCl, the ratio *P*₃₅:*P*₃₇ = 3.01, while for exposed *trans*-ClHC=CHCl *P*₃₅:*P*₃₇ = 2.73. A small but significant enrichment of ³⁷Cl in the ex-




Figure 3. Mass spectrum of the m/e 35 and 37 peaks for (a) naturally occurring $trans$ - $ClHC=CHCl$ and (b) $trans$ - $ClHC=CHCl$ separated from the exposed reaction mixture. The $\times 3$ peak is recorded on a scale three times smaller than the $\times 1$ peak.

posed starting material is observed, demonstrating laser-controlled isotope interchange. Further confirmation of enrichment is provided by examining the ratio of other mass peaks. For example, $P_{48}:P_{50} = 4.26$ for reference $trans$ - $ClHC=CHCl$ while $P_{48}:P_{50} = 3.68$ for exposed $trans$ - $ClHC=CHCl$. Here P_{48} is mainly due to $^{12}CH^{35}Cl$ and P_{50} to $^{12}CH^{37}Cl$.

The cis - $ClHC=CHCl$ photoproduct also shows isotope enrichment, e.g., $P_{35}:P_{37} = 2.89$. In addition to the gaseous products, a pink nonvolatile liquid is formed during irradiation which has been identified as the photoaddition product $C_2H_2ICl_3$.

So far, our most impressive isotope separation has been achieved using 1,2-dibromoethylene as a scavenger under the same condition as $trans$ -dichloroethylene. Several photoproducts are formed, viz., C_2H_2BrCl and cis - and $trans$ - $ClHC=CHCl$, presumably resulting from the subsequent reaction of C_2H_2BrCl with ICl^* . For the $trans$ - $ClHC=CHCl$ product, the $P_{35}:P_{37}$ ratio changes from 3.01 to 2.03, as shown in Figure 3. Moreover, the ICl in the exposed mixture is found to be enriched in ^{35}Cl , since its ^{37}Cl has been depleted by reaction.

Various laser isotope separation schemes have been proposed, and in some cases demonstrated.²⁻¹¹ In particular, photochemical enrichment of chlorine isotopes has been previously reported by Lamotte, Dewey, Keller, and Ritter¹¹ who used the photoaddition reaction between electronically excited thiophosgene and diethoxyethylene. By exciting the individual rotational lines of isotopically selected $CSCl_2$, they obtained an enrichment in the unreacted $CSCl_2$ comparable to ours. We anticipate, however, that our degree of isotopic enrichment will increase markedly when our laser line width is narrowed so as to excite individual rotational lines of $I^{35}Cl$ or $I^{37}Cl$. Moreover, it should be stressed here that we now have a convenient system for the study of the dynamics of organic gas-phase photochemical reactions in which mechanisms for isotopic scrambling and retention—a central problem in isotope separation schemes—can be well understood.

Acknowledgment. This work is dedicated to the memory of the late Professor T. I. Taylor (Department of Chemistry, Columbia University). In this work, we used the mass spectrometer which he constructed. Financial support was provided by the National Science Foundation.

References and Notes

- (1) (a) E. Hulthén, N. Johansson, and U. Pilsäter, *Ark. Fys.*, **14**, 31 (1958); (b) E. Hulthén, N. Järlsäter, and L. Koffman, *ibid.*, **18**, 479 (1960).
- (2) W. B. Tiffany, H. W. Moos, and A. L. Schawlow, *Science*, **157**, 40 (1967); W. B. Tiffany, *J. Chem. Phys.*, **48**, 3019 (1968).
- (3) R. L. Farrar, Jr., and D. F. Smith, "Photochemical Isotope Separation as Applied to Uranium", Report K-L-3054, Rev. 1 (March 15, 1972) Union Carbide Corporation, Oak Ridge, Tenn.
- (4) V. S. Letokhov, *Science*, **180**, 451 (1973).
- (5) C. B. Moore, *Acc. Chem. Res.*, **6**, 323 (1973).
- (6) E. S. Yeung and C. B. Moore, *Appl. Phys. Lett.*, **21**, 109 (1971).
- (7) R. V. Ambartsumyan, V. S. Letokhov, G. N. Makarov, and A. A. Pureskii, *Zh. Eksp. Teor. Fiz. Pis'ma Red.*, **17**, 91 (1973).
- (8) Post deadline papers at the VIII International Conference on Quantum Electronics, San Francisco, Calif., June 1974, reported the laser isotope separation of: (a) uranium by S. A. Tuccio, J. W. Dubrin, O. G. Peterson, and B. B. Snavely; (b) barium by A. Bernhardt, D. Deurre, J. Simpson, and L. Wood; (c) boron by S. Rockwood and S. W. Rabideau; (d) calcium by U. Brinkmann, W. Hartig, H. Telle, and H. Walther.
- (9) S. R. Leone and C. B. Moore, *Phys. Rev. Lett.*, **33**, 269 (1974).
- (10) S. M. Freund and J. J. Ritter, *Chem. Phys. Lett.*, submitted for publication.
- (11) M. Lamotte, H. J. Dewey, R. A. Keller, and J. J. Ritter, *Chem. Phys. Lett.*, **30**, 165 (1975).

Dirkson D.-S. Liu, Saswati Datta, Richard N. Zare*


Department of Chemistry, Columbia University
New York, New York 10027

Received January 27, 1975

9-BBN Ate Complexes as a New Type of Reducing Agent for the Selective Reduction of Tertiary Alkyl, Benzyl, and Allyl Halides to Hydrocarbons

Sir:

We wish to report on the interesting reducing character of 9-borabicyclo[3.3.1]nonane (9-BBN) ate complexes (**1**). The reagents enable the selective removal of tertiary alkyl, benzyl, and allyl halides to afford the corresponding hydrocarbons in excellent yields without concomitant attack on secondary, primary, and aryl derivatives (eq 1).



Although hydride character in the tetraalkylboron compounds was originally proposed by Wittig,¹ only sparse reports of their reducing ability have appeared.² In the course of studies on borate complexes, certain observations suggested that the bridgehead hydrogens of **1** must be exceptionally labile as hydride sources. Accordingly, we examined the reaction of benzyl chloride with the n -butyl ate complexes³ of tri- n -butylborane, tri- sec -butylborane, and B - n -Bu-9-BBN. As is apparent from Table I, the secondary

Table I. Reactions of Benzyl Chloride with Ate Complexes^a

Ate complex	Toluene (%) ^b	Benzyl chloride (%) ^b
Li (n -Bu) ₃ B	0	100
Li (n -Bu)(sec -Bu) ₂ B	75	18
1a	100	0

^a Benzyl chloride (10 mmol) was added to the mixture of ate complex (10 mmol) and n -hexane at 0°. The resultant mixture was stirred at 20° for 3 hr, and then oxidized with $NaOH-H_2O_2$. ^b By GLPC analysis.